



On lower bounds for the largest eigenvalue of a symmetric matrix

S.G. Walker^{a,*}, P. Van Mieghem^b

^a *Institute of Mathematics, Statistics and Actuarial Science, University of Kent, Canterbury CT2 7NZ, UK*

^b *Delft University of Technology, Faculty of Electrical Engineering, Mathematics and Computer Science, P.O. Box 5031, 2600 GA Delft, The Netherlands*

Received 17 September 2007; accepted 13 March 2008

Available online 22 April 2008

Submitted by S. Friedland

Abstract

We consider lower bounds for the largest eigenvalue of a symmetric matrix. In particular we extend a recent approach by Piet Van Mieghem.

© 2008 Elsevier Inc. All rights reserved.

AMS classification: Primary 15A42; Secondary 30B10

Keywords: Largest eigenvalue; Positive definite matrix; Symmetric matrix

1. Introduction

Let $\lambda_{\max}(A)$ be the largest eigenvalue of a symmetric $m \times m$ matrix $A = (a_{ij})$. Since

$$\lambda_{\max}(A) = \max_{x \neq 0} \frac{x^T A x}{x^T x}$$

it clearly follows that a lower bound for $\lambda_{\max}(A)$ is given by

$$\lambda_{\max}(A) \geq \frac{u^T A u}{u^T u} \tag{1}$$

* Corresponding author. Tel.: +44 (0) 1227 823800; fax: +44 (0) 1227 827932.

E-mail addresses: S.G.Walker@kent.ac.uk (S.G. Walker), P.F.A.VanMieghem@tudelft.nl (P. Van Mieghem).

where $u^T = (1 \cdots 1)$. Note that

$$N_1 = u^T A u = \sum_{ij} a_{ij},$$

$u^T u = m$ and N_1/m is a commonly used lower bound for $\lambda_{\max}(A)$. Recent work on lower bounds for a symmetric matrix has been done by Van Mieghem [2]. He showed that

$$\lambda_{\max}(A) \geq \frac{N_1}{m} + 2 \left(\frac{N_3}{2m} - \frac{N_1 N_2}{m^2} + \frac{N_1^3}{2m^3} \right) \lambda_0^{-2} + O(t^{-4}), \tag{2}$$

where $t \geq T$, $\lambda_0 = t\sqrt{m}$,

$$T = \frac{1}{\sqrt{m}} \max_{1 \leq j \leq m} \left(a_{jj} + \sum_{i \neq j} |a_{ij}| \right), \tag{3}$$

and $N_k = u^T A^k u$ with $N_0 = m$.

The aim in the current paper is to extend the results of Van Mieghem [2]. The central idea of the paper is to apply the classic bound to transforms of A . Applying standard bounds to transformed matrices which result in improved bounds has recently been exploited in Walker [3,4] and Liu et al. [1]. We derive the general lower bound in Section 2, where we also consider some specific cases. Section 3 provides a further useful result when A is positive definite and finally Section 4 concludes with a numerical example.

2. Lower bounds for symmetric matrices

Consider the $m \times m$ symmetric matrix

$$A_t = \sum_{k=0}^{\infty} f_k A^k t^{-k},$$

where the Taylor series $\sum_{k=0}^{\infty} f_k x^k = f(x)$ converges for $|x| < R_f$, where $R_f > 0$ is the radius of convergence. If λ is an eigenvalue of A , corresponding to eigenvector v , then

$$A_t v = \sum_{k=0}^{\infty} f_k A^k t^{-k} v = \sum_{k=0}^{\infty} f_k \lambda^k t^{-k} v = f\left(\frac{\lambda}{t}\right) v.$$

The series converges for any eigenvalue of A provided we choose $t > \tilde{\lambda}/R_f$, where $\tilde{\lambda} = \max_{1 \leq j \leq m} \{|\lambda_j|\}$.

If $f(x)$ is real for real x and increasing, then $\lambda_{\max}(A_t) = f\left(\frac{\lambda_{\max}(A)}{t}\right)$. Next, we apply the classical bound (1) to A_t and obtain

$$\lambda_{\max}(A_t) \geq \frac{u^T A_t u}{m} = \frac{1}{m} \sum_{k=0}^{\infty} f_k (u^T A^k u) t^{-k} = \frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k}.$$

It follows from (1) that $N_k \leq m \lambda_{\max}(A^k)$. Since $\lambda_{\max}(A^k) \leq \tilde{\lambda}^k$, we have that $N_k \leq m \tilde{\lambda}^k$ and this inequality shows that the series $\sum_{k=0}^{\infty} f_k N_k t^{-k}$ indeed converges for $t > \tilde{\lambda}/R_f$.

Since also the inverse function $f^{-1}(x)$ is increasing when $f(x)$ is increasing such that

$$\lambda_{\max}(A) = t f^{-1}(\lambda_{\max}(A_t)),$$

we arrive at the inequality

$$\lambda_{\max}(A) \geq t f^{-1} \left(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k} \right). \tag{4}$$

The best possible bound is reached when the right hand side in (4) is optimized over all increasing functions f . Obviously the set of increasing functions includes the case $f(x) = x$ and for this increasing function we obtain the classic inequality $\lambda_{\max}(A) \geq N_1/m$. Hence (4) is at least as good as the classic bound when optimized over all increasing functions. In fact as we will see in Section 3, when A is positive definite, it turns out that the worst f is indeed $f(x) = x$.

The function $f^{-1} \left(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k z^k \right)$ is expanded in a series around $z = 1/t = 0$ in Appendix A to obtain

$$\lambda_{\max}(A) \geq t f^{-1} \left(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k} \right) = \sum_{k=1}^{\infty} c_k t^{1-k} \tag{5}$$

and the general term c_k is given in (8). Explicitly, the first few coefficients c_k are

$$\begin{aligned} c_1 &= \frac{N_1}{m} \\ c_2 &= \frac{f_2}{f_1} \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) \\ c_3 &= \frac{f_3}{f_1} \left(\frac{N_3}{m} - \frac{N_1^3}{m^3} \right) + \frac{2f_2^2}{f_1^2} \left(\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} \right) \\ c_4 &= \frac{f_4}{f_1} \left(\frac{N_4}{m} - \frac{N_1^4}{m^4} \right) + \frac{f_2 f_3}{f_1^2} \left(\frac{5N_1^4}{m^4} - \frac{3N_1^2 N_2}{m^3} - \frac{2N_1 N_3}{m^2} \right) \\ &\quad + \frac{f_2^3}{f_1^3} \left(-\frac{5N_1^4}{m^4} + \frac{6N_1^2 N_2}{m^3} - \frac{N_2^2}{m^2} \right). \end{aligned}$$

If $R_{f^{-1}}$ is the radius of convergence of the Taylor series of $f^{-1}(x)$ around f_0 , then

$$f^{-1} \left(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k} \right) = f^{-1} \left(f_0 + \frac{1}{m} \sum_{k=1}^{\infty} f_k N_k t^{-k} \right)$$

indicates that convergence requires that $\frac{1}{m} \sum_{k=1}^{\infty} f_k N_k t^{-k} < R_{f^{-1}}$. Using $N_k \leq m \tilde{\lambda}^k$, the series is bounded by

$$\frac{1}{m} \sum_{k=1}^{\infty} f_k N_k t^{-k} \leq \sum_{k=1}^{\infty} f_k \tilde{\lambda}^k t^{-k} = f \left(\frac{\tilde{\lambda}}{t} \right) - f_0$$

from which $f \left(\frac{\tilde{\lambda}}{t} \right) < f_0 + R_{f^{-1}}$ and thus, that $t > \frac{\tilde{\lambda}}{f^{-1}(f_0 + R_{f^{-1}})}$. Combined with the above bounds on t , convergence of $\sum_{k=1}^{\infty} c_k t^{1-k}$ requires that

$$t > \tilde{\lambda} \max \left(\frac{1}{R_f}, \frac{1}{f^{-1}(f_0 + R_{f^{-1}})} \right) \tag{6}$$

and, in practice, $t > \tilde{T}\sqrt{m} \max\left(\frac{1}{R_f}, \frac{1}{f^{-1}(f_0+R_{f-1})}\right)$, where

$$\tilde{T}\sqrt{m} = \max_{1 \leq j \leq m} \left\{ \sum_{i=1}^m |a_{ij}| \right\},$$

since it is well known that $\tilde{\lambda} < \tilde{T}\sqrt{m}$.

2.1. Examples

If $f_k = 1$, then $f(x) = \frac{1}{1-x}$ and $f^{-1}(x) = 1 - \frac{1}{x}$. The Taylor series of $f(x)$ around $x = 0$ has $R_f = 1$, while the Taylor series of $f^{-1}(x)$ around $f(0) = 1$ has radius of convergence $R_{f^{-1}} = 1$. Hence, the bound (6) for t yields $t > 2\tilde{T}\sqrt{m}$ and we find from (5)

$$\lambda_{\max}(A) \geq \frac{N_1}{m} + \frac{1}{t} \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) + 2 \left(\frac{N_3}{2m} - \frac{N_1 N_2}{m^2} + \frac{N_1^3}{2m^3} \right) \frac{1}{t^2} + O(t^{-3}). \tag{7}$$

The bound (7) is very similar to the Van Mieghem [2] expression (2), except we have an additional $1/t$ term which is positive. Note that the $1/t^2$ term is not necessarily positive. On the other hand, the bound on t in (2) is less than half as large as $2\tilde{T}\sqrt{m}$ here.

If we choose $f(x) = (1-x)^\alpha$, then the Taylor coefficients around $x = 0$ are $f_k = (-1)^k \binom{\alpha}{k}$ and $R_f = 1$. The inverse function $f^{-1}(x) = 1 - x^{\frac{1}{\alpha}}$ has a radius of convergence around $f(0) = 1$ equal to $R_{f^{-1}} = 1$. Using (6), we have that $t > \tilde{T}\sqrt{m} \max\left(1, \frac{1}{1-2^{\frac{1}{\alpha}}}\right)$. For $\alpha = -|\beta| < 0$, where $f_k = \binom{|\beta|-1+k}{k}$ and $t > \frac{\tilde{T}\sqrt{m}}{1-2^{-\frac{1}{|\beta|}}}$, the lower bound (5) up to $O(t^{-3})$ is

$$\begin{aligned} \lambda_{\max}(A) \geq & \frac{N_1}{m} + \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) \frac{(|\beta|+1)}{2t} \\ & + \left\{ \frac{(|\beta|+2)}{3(|\beta|+1)} \left(\frac{N_3}{m} - \frac{N_1^3}{m^3} \right) + \left(\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} \right) \right\} \frac{(|\beta|+1)^2}{2t^2}. \end{aligned}$$

To compare with (7) where $|\beta| = 1$, we write $t = t_1 \frac{1}{2(1-2^{-\frac{1}{|\beta|}})}$, where $t_1 > 2\tilde{T}\sqrt{m}$,

$$\begin{aligned} \lambda_{\max}(A) \geq & \frac{N_1}{m} + \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) \frac{(|\beta|+1) \left(1 - 2^{-\frac{1}{|\beta|}}\right)}{t_1} \\ & + \left\{ \frac{(|\beta|+2)}{3(|\beta|+1)} \left(\frac{N_3}{m} - \frac{N_1^3}{m^3} \right) + \left(\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} \right) \right\} \frac{2(|\beta|+1)^2 \left(1 - 2^{-\frac{1}{|\beta|}}\right)^2}{t_1^2} \end{aligned}$$

This shows that the coefficient of $\frac{1}{t_1}$ is larger than in the $\beta = 1$ case provided $|\beta| < 1$. In that case, however, the coefficient of $\frac{1}{t_1^2}$ has a smaller positive $\frac{(|\beta|+2)}{3(|\beta|+1)}$ factor. The argument shows that, depending on the values of N_k , we may fine-tune β to produce a larger lower bound.

Finally, consider $f(x) = e^{ax}$ for which $f_k = \frac{a^k}{k!}$ and $R_f \rightarrow \infty$. The inverse function $f^{-1}(x) = \frac{1}{a} \log x$ has a Taylor series around $f(0) = 1$ with $R_{f^{-1}} = 1$. The bound (6) becomes $t > \frac{a\tilde{T}\sqrt{m}}{\log 2}$ and (5) up to $O(t^{-3})$ is

$$\lambda_{\max}(A) \geq \frac{N_1}{m} + \frac{1}{2} \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) \frac{a}{t} + \left\{ \frac{a}{3} \left(\frac{N_3}{m} - \frac{N_1^3}{m^3} \right) + \left(\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} \right) \right\} \frac{a^2}{2t^2}$$

Comparison with (7) via $t = \frac{at_1}{2 \log 2}$ gives

$$\lambda_{\max}(A) \geq \frac{N_1}{m} + \left(\frac{N_2}{m} - \frac{N_1^2}{m^2} \right) \frac{\log 2}{t_1} + \left\{ \frac{a}{3} \left(\frac{N_3}{m} - \frac{N_1^3}{m^3} \right) + \left(\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} \right) \right\} \frac{2 \log^2 2}{t_1^2}.$$

The coefficient of $\frac{1}{t_1}$ is now smaller than in the $\beta = 1$ case, but the value of a can be freely chosen in the coefficient of $\frac{1}{t_1^2}$ (ignoring higher order terms).

Another possible sequence of functions to consider is $f(x) = x^k$ for odd k . As has been mentioned the case $k = 1$ provides the classic bound. For these functions the inverse is trivial and hence bounds are easily available.

3. Positive definite case

When A is positive definite we have the following key result:

Lemma 3.1. *It is that $N_k \geq N_1^k / m^{k-1}$ for all $k = 1, 2, \dots$*

Proof. It is well known that we can write

$$A = QDQ^T = \sum_{j=1}^m \lambda_j^k v_j v_j^T,$$

where Q is an orthogonal matrix with column eigenvectors $\{v_j\}$, and D is a diagonal matrix with entries the eigenvalues $\{\lambda_j\}$. So

$$N_k = \sum_{j=1}^m \lambda_j^k u^T v_j v_j^T u$$

and $u^T v_j v_j^T u = (u^T v_j)^2$ with

$$\sum_{j=1}^m (u^T v_j)^2 = m.$$

Hence, $N_k = E(A^k)$ with $P(A = \lambda_j) = (u^T v_j)^2 / m$; and, since $\lambda_j > 0 \forall j$, a consequence of A being positive definite, it is that $A > 0$ with probability one, and using Jensen’s inequality, it is that $E(A^k) \geq \{E(A)\}^k$. So $N_k = mE(A^k) \geq m\{E(A)\}^k = N_1^k / m^{k-1}$, completing the proof. \square

Applying Lemma 3.1 shows that

$$\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k} \geq \sum_{k=0}^{\infty} f_k \frac{N_1^k}{m^k} t^{-k} = f \left(\frac{N_1}{tm} \right).$$

Hence, the inequality (4) is lower bounded by

$$\lambda_{\max}(A) \geq t f^{-1} \left(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k t^{-k} \right) \geq t f^{-1} \left(f \left(\frac{N_1}{tm} \right) \right) = \frac{N_1}{m}.$$

In other words, if A is symmetric and positive definite and if $f(x)$ is increasing, then (4) is at least as sharp as the classical bound N_1/m .

A better bound is achieved when all c_k in (5) are made larger than those in (7). This seems possible, because Lemma 3.1 states that the prefactor $N_k/m - (N_1/m)^k$ of f_k/f_1 in (8) is always positive. For, choose $f_2 > 1$, then c_2 is larger. However, increasing f_2 has a negative effect on c_3 since $\frac{N_1^3}{m^3} - \frac{N_1 N_2}{m^2} < 0$. This effect can be compensated by choosing f_3 sufficiently large. A same argument applies for all other terms: there is always the possibility to choose in c_k the highest Taylor coefficient f_k , that is multiplied by $\left(\frac{N_k}{m} - \frac{N_1^k}{m^k}\right) > 0$, sufficiently large to compensate for the possible decrease in c_k by augmenting lower order Taylor coefficients f_j with $j < k$. It is a matter of optimizing the Taylor coefficients f_k and the bound (6) on t .

4. Numerical examples

Here we consider a specific example when

$$A = \begin{pmatrix} -1 & \sqrt{6} \\ \sqrt{6} & -2 \end{pmatrix}.$$

The eigenvalues of A are 1 and -4 and we have $N_1 = 1.8990$, $N_2 = 2.3031$ and $N_3 = 0.6867$. Hence, the classic bound is given by $N_1/m = 0.9495$. On the other hand, using (7) with $t = 2\tilde{T}\sqrt{m} = 8.8990$, we obtain a lower bound for $\lambda_{\max}(A)$ as 0.9521, which obviously improves on 0.9495.

Now we consider the example when A is a 10×10 symmetric matrix and for $j = 1, \dots, i$ we have $a(i, j) = 2j - i$. Then we have $N_1 = 55$, $N_2 = 3553$ and $N_3 = 108823$. Hence the classic bound is given by 5.5. The bound (7) with $t = 2\tilde{T}\sqrt{10}$, and $\tilde{T}\sqrt{10} = 50$, is given by the improved lower bound of 9.465. However, for this example, the function $f(x) = x^3$ provides the lower bound of $(N_3/10)^{1/3} = 22.16$.

If we now take $a(i, j) = 2j - 3i$, $j \leq i$, and A is again a 10×10 symmetric matrix, then $N_1 = -1375$, $N_2 = 194425$ and $N_3 = -27325375$. Also $\tilde{T}\sqrt{10} = 190$. So the classic bound is -137.5 and the bound (7) with $t = 2\tilde{T}\sqrt{10}$ is -136.00 . On this occasion the bound based on $f(x) = x^3$ is given by -139.8 , which is smaller than the classic bound. The bound (2) is given by -137.00 which improves on the classic bound but is worse than (7).

Appendix A. Taylor expansion of $f^{-1}(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k z^k)$ around $z = 0$

We now expand $f^{-1}(\frac{1}{m} \sum_{k=0}^{\infty} f_k N_k z^k)$ in a series around $z = \frac{1}{t} = 0$ by invoking characteristic coefficients, defined e.g. in [2, Appendix]. We apply the general expansion (deduced from [2, Appendix]), provided that $f(z_0) = h(z_0)$

$$\begin{aligned}
 f^{-1}(h(z)) &= z_0 + \sum_{m=1}^{\infty} \frac{h_m(z_0)}{f_1(z_0)} (z - z_0)^m \\
 &+ \sum_{m=2}^{\infty} \sum_{n=2}^m \left(\sum_{k=1}^{n-1} (-1)^k \binom{n+k-1}{k} \right) f_1^{-n-k}(z_0) s^*[k, n-1]|_{f(z)}(z_0) \\
 &\times \frac{s[n, m]|_{h(z)}(z_0)}{n} (z - z_0)^m
 \end{aligned}$$

to $h(z) = \frac{1}{m} \sum_{k=0}^{\infty} f_k N_k z^k = f_0 + \frac{f_1 N_1}{m} z + \frac{1}{m} \sum_{k=2}^{\infty} f_k N_k z^k$ and $z_0 = 0$. Then,

$$\begin{aligned}
 f^{-1}(h(z)) &= \frac{1}{m f_1} \sum_{k=1}^{\infty} f_k N_k z^k \\
 &+ \sum_{m=2}^{\infty} \sum_{n=2}^m \left(\sum_{k=1}^{n-1} (-1)^k \binom{n+k-1}{k} \right) f_1^{-n-k} s^*[k, n-1]|_{f(z)} \frac{s[n, m]|_{h(z)}}{n} z^m.
 \end{aligned}$$

Hence,

$$c_k = \frac{f_k N_k}{m f_1} + \sum_{n=2}^k \left(\sum_{j=1}^{n-1} (-1)^j \binom{n+j-1}{j} \right) f_1^{-n-j} s^*[j, n-1]|_{f(z)} \frac{s[n, k]|_{h(z)}}{n}.$$

This can be simplified using $s[k, k] = f_1^k$ and $s[1, m] = f_m$ to explicitly obtain the prefactor of the highest Taylor coefficient f_k in c_k ,

$$\begin{aligned}
 c_k &= \frac{f_k N_k}{m f_1} + \frac{1}{k} \left(\frac{N_1}{m} \right)^k \sum_{j=1}^{k-1} (-1)^j \binom{k+j-1}{j} f_1^{-j} s^*[j, k-1]|_{f(z)} \\
 &+ \sum_{n=2}^{k-1} \left(\sum_{j=1}^{n-1} (-1)^j \binom{n+j-1}{j} \right) f_1^{-n-j} s^*[j, n-1]|_{f(z)} \frac{s[n, k]|_{h(z)}}{n}
 \end{aligned}$$

or

$$\begin{aligned}
 c_k &= \frac{f_k}{f_1} \left(\frac{N_k}{m} - \left(\frac{N_1}{m} \right)^k \right) + \frac{1}{k} \left(\frac{N_1}{m} \right)^k \sum_{j=2}^{k-1} (-1)^j \binom{k+j-1}{j} f_1^{-j} s^*[j, k-1]|_{f(z)} \\
 &+ \sum_{n=2}^{k-1} \left(\sum_{j=1}^{n-1} (-1)^j \binom{n+j-1}{j} \right) f_1^{-n-j} s^*[j, n-1]|_{f(z)} \frac{s[n, k]|_{h(z)}}{n}. \tag{8}
 \end{aligned}$$

References

[1] X-Q. Liu, T-Z. Huang, Y-D. Fu, A note on improvements on bounds for nonmaximal eigenvalues of symmetric positive matrices, *Linear Algebra Appl.* 419 (2006) 612–617.
 [2] P. Van Mieghem, A new type of lower bound for the largest eigenvalue of a symmetric matrix, *Linear Algebra Appl.* 427 (2007) 119–129.

- [3] S.G. Walker, On recent Cheeger type bounds for non-maximal eigenvalues applied to positive matrices, *SIAM J. Matrix Anal. Appl.* 25 (2003) 574–581.
- [4] S.G. Walker, Improving bounds for nonmaximal eigenvalues of nonnegative matrices, *Linear Algebra Appl.* 397 (2005) 133–139.